

02/04/02

JC698 U.S. PTO

NEW SAFE BOTANICAL DRUG FOR TREATING MALIGNANT PLEURAL  
EFFUSION AND CANCER AND INCREASING IMMUNE FUNCTION

JC903 U.S. PTO  
10/062316  
02/04/02

**Background of the Invention**

This invention relates to new safe botanical drug, which is treatment and prevention of malignant pleural effusion and cancer and increase immune function. Specifically, this invention provides the safe botanical drug comprising Polysaccharide of Dang Gui (PDG) and Lan Xiang Xi (LX). New botanical drug, which treats and prevents cancer through killing cancer cells and increasing immune function at same time, is safe.

**Description of Prior Art**

Cancer is the second leading cause of death in the United States, and the incidence of cancer continues to climb annually. In recent years, about 1 million new cases of cancer are diagnosed yearly in the U.S. About half million of people and 7 million people of annual died in the US and in the world, respectively. A lot of anticancer drugs including chemical and antibiotics have effects to kill cancer cells. But it also kills off some normal human cells, appears many kinds of side effects, among them the inhibition of bone marrow and tract are the most common. For example, Cyclophosphamide is a chemotherapeutic drug, which is highly effective against a wide range of human cancers. Unfortunately, it does damage to hemocytogenesis organs, alimentary tract and decreasing immune function. Adriamycin, a sensitive anticancer antibiotic, has seriously cardiotoxicity. Therefore, decreasing toxicity of anticancer drugs is very important.

Many reports indicated that the side effects of plant's anticancer drugs are lower than chemical and antibiotic's anticancer drugs. Therefore, the development of plant drug has progressed very fast now. Taxol, for example, is a novel anticancer plant drug isolated from the needles and bark of the western yew, *Taxus brevifolia*. It is the prototype for a new class of antitumor drugs, which are characterized by their capacity to promote the assembly of microtubules. Clinical trials conducted in the late 1980s and early 1990s demonstrated impressive clinical activities against advanced ovarian and breast cancer. Taxol still has some side effects, for example, myelosuppression, diarrhea, emesis, oligospermia, cellular depletion in lymphoid tissues, changes in serum hepatic enzymes, and elevations in cholesterol and triglycerides were observed. More important, taxol has two big problems. The first problem is that natural source of taxol is very

limited. And the second problem is that taxol is a poor-water soluble. Vehicles for parental administration on taxol cause serious side effects.

Recently a number of gene expression systems have been developed that can be regulated by the administration of specific small molecule drugs. It is important that the small molecule drugs are easily administered. Some of the earliest systems include exposure to heavy metal, and steroid hormones, but they are not suited for in vivo human. These systems, however, are suitable to study for the effect of small molecular drugs on controllable oncogenes, and to study for the treatment of cancer.

So far, no one drug has been succeeded to treat or prevent cancer by control cancer cells and without adverse side effects.

### **Summary of the Invention**

This invention relates to new safe botanical drug, which is treatment and prevention of treating malignant pleural effusion, increase immune function and treating cancer. Specifically, this invention provides the safe botanical drug comprising polysaccharide of Dang Gui (PDG) and Lan Xiang Xi (LX).

New botanical drug, which treats and prevents cancer through killing or controlling cancer cells and increasing immune function in same time, is safe.

Methods of killing or controlling cancer cells include inhibiting oncogenes, increasing activity of tumor suppressor, inducing differentiation of cancer cells, inhibiting cancer cells proliferation, inducing apoptosis of cancer cells, inhibiting growth of transplanted cancer and inhibiting cancer incidence.

### **Detailed Description of the Invention**

Recent developments in cancer research, more specifically, in molecular biology, biochemical aspects, indicated that controlling cancer cells is more important than killing cancer cells. For example, control oncogene expression, apoptosis, and differentiation of cancer cells are several key steps in process of control cancer. The control of oncogene expression, apoptosis and differentiation of cancer is focus in cancer biology and molecular pharmacology of anticancer drug. For example, concentration of oncogene c-myc and c-myb RNA is rapid increased in human leukemia and other cancer cell, and a rapid decline in the expression of c-myc and c-myb

RNA was seen in cancer cells induced to differentiate. To inhibit oncogene expression, apoptosis and differentiation of cancer cells can control and destroy carcinogenesis and stop cancer progression.

It is more important that LX is successfully used in clinic for malignant pleural effusion and treatment cancer. For example, at least have 600 cases clinic reported that LX treated patients with malignant pleural effusion higher than 70%. Also, LX could effective treat lung cancer and increasing immune function.

The following specific examples will provide detailed illustrations of methods of producing relative drugs, according to the present invention and pharmaceutical dosage units containing relative drugs. Moreover, examples described pharmaceutical characters of drugs, which demonstrated its effectiveness in control of cancer cells. These examples are not intended, however, to limit or restrict the scope of the invention in any way, and should not be construed as providing conditions, parameters, reagents, or starting materials which must be utilized exclusively in order to practice the present invention.

#### **Example 1 Extraction of Polysaccharides of Dang Gui (PDG)**

2,000 ml of water was added to 1,000 g of dry powder of Dang Gui, a Chinese herb. The mixture was heated to boil and simmered for one and one-half hours after boiling. This water extraction was repeated once and the two extracts were combined and filtered. The filtrate was concentrated under reduced pressure to approximately 500 ml and 95% ethanol was added to the concentrate to yield a final alcohol concentration of 60%. The resulting solution was filtered to recover a precipitate which was dissolved in an appropriate amount of water, the resulting solution was filtered to remove residue and a filtrate was saved. The filtrate was concentrated under reduced pressure to 200 ml and 95% ethanol added to the concentrate to yield a final alcohol concentration of 80%. The solution was then discarded and the precipitate was washed three times with 95% ethanol and then twice with acetone and ether consecutively. The product was vacuum dried, and the resulting powder was polysaccharides of Dang Gui.

#### **Example 2 LX extraction**

One kg of plant powder was extracted 5 L of water at room temperature for 12 hours. The powder of plant named *Dryobalanops aromatica* Gaerin or Wen E Shu was recovered by filtration. Filtrate

A was saved and the powder filtercake was extracted with 4 L of water at room temperature for 10 hours. The mixture was filtered. Filtrate B was saved and powder filtercake was extracted for 3 L of water at room temperature for 8 hours. The mixture was filtered and filtrate C was saved. Filtrate A, B, and C was combined and distilled under reduced pressure for 32 hours. The distilled mixture was separated. The oil fraction was saved and kept at temperature at 0°C. The oil distilled under reduced pressure, (50° – 80°C/40 Pa) and fraction A was collected. Fraction A distilled under reduced pressure (76°-78°C/40 Pa) and fraction B was collected. Fraction B was then chromatographed on silica gel G, using petroleum ether as developing solvent. The solvent collected and dried. The final product is LX.

LX has the following chemistry data.

Molecular form:  $C_{15}H_{24}$

Molecular weight. 204

Mp: 114~118°C

$[\alpha]^{16}_D$  : -15°

IR $\nu^{KBr}$   $cm^{-1}$ : 3090, 2975, 2860, 1642, 1440, 1375, 1002, 910, 888 (  $C=CH_2$ );

PMR ( $CCl_4$ )  $\delta$ : 0.97 (3H, s), 1.7 (6H, s), 4.4~5.6 (6H, m), 5.7 (1H, dd, =  $OH_2$ );

MSm/e (%): 204 ( $M^+$ ), 147 (33), 121 (41), 107 (54), 93 (89), 81 (100), 79 (44), 68 (74), 67 (52), 55 (41), 53 (33), 41 (52).

### **Example 3 Injection solution of LX**

Four (4) volumes of 95% ethanol were added to LX. The solution was allowed to stand for 24 hours and then filtered. The filtrate distilled under reduced pressure and the ethanol recovered. Six volumes of 95% ethanol were added to the residue. After standing for another 24 hours, the solution was filtered. The filtrate distilled under reduced pressure and ethanol recovered. The residue was then distilled until there was no remaining smell of alcohol. Sufficient distilled water was added to dissolve the residue, and the solution was filtered to remove any undissolved material. Pharmaceutical glycerin was added to the solution. The solution was then fine filtered, and the volume adjusted to 500 liters with distilled water. After additional fine filtering, the solution was sealed in 2ml sterile ampoules, which were further sterilized and sealed. Each ampoule contained 5 mg of LX per milliliter of solution.

#### **Example 4 Preparation of LX-containing sterically stabilized liposomes (LX-SSL)**

Hydrogenated phosphatidyl choline (PC), phosphatidyl glycerol (PG), and phosphatidyl serine (PS) were extracted from soybean. All above lipids were finally purified on silicic acid columns, shown to be pure by thin-layer chromatography and stored in chloroform in sealed ampules under nitrogen until use. Phospholipids mixed with cholesterol (CHOL) and long-chain alcohol. The solvent was removed under reduced pressure by a rotary evaporator. The lipids were then purged with nitrogen. Lipids were redissolved in the organic phase and reversed phase will be formed. LX-containing solution was added at these lipid systems, and resulting two-phase system was sonic 3 minutes until the mixture homogeneous that did not separate for at least two hours. A typical preparation contained  $3.3 \times 10^{-3}$  M of phospholipid and  $3.3 \times 10^{-3}$  M of cholesterol in 1 litre of phosphate-buffered saline and 3 liters of solvent. LX-SSL were sealed and sterilized. The size of the vesicles was determined by a dynamic light scattering technique. When PG/PC/CHOL were 1:4:5, diameter of liposomes was 20-50 nm (range). LX-SSL was very stabilized in at least nine months.

So far, many articles reported drug-containing liposomes. However, liposomes are not stabilized enough. Therefore it is difficult to be used for pharmaceutical industry. In accordance with this invention, LX-SSL is very stabilized in at least nine months. Therefore LX-SSL can be used in industry. LX-SSL can enhance targeting and improve pharmaceutical activity of LX. The results show that LX-SSL is safer than LX. LX + PDG-SSL is safer than LX + PDG. Also it is very important that all lipids are extracted from soybean. Therefore it is very safe for human being. In fact, many articles reported lipids, which used for drug-containing liposomes, are syntheses by organic chemistry. However synthetic lipids have some side effects, therefore, according to this invention, methods of preparing of LX-SSL and phospholipids, which extracted from soybean, are very safe.

#### **Example 5 Preparation of botanical drug**

1 kg polysaccharide of Dang Gui and 1 kg of LX produced in accordance with the preceding example, were thoroughly mixed and agitated until a homogeneous mixture of the two components was obtained. Dosage form was capsule or tablet, which includes in addition pharmaceutically acceptable binders and excipients.

#### **Example 6 Preparation of PDG-containing sterically stabilized liposomes (PDG-SSL)**

Hydrogenated phosphatidyl choline (PC), phosphatidyl glycerol (PG), and phosphatidyl serine (PS) were extracted from soybean. All above lipids were finally purified on silicic acid columns, shown to be pure by thin-layer chromatography and stored in chloroform in sealed ampoules under nitrogen until use. Phospholipids mixed with cholesterol (CHOL) and long-chain alcohol. The solvent was removed under reduced pressure by a rotary evaporator. The lipids were then purged with nitrogen. Lipids were redissolved in the organic phase and reversed phase will be formed. PDG-containing solution was added at these lipid systems, and resulting two-phase system was sonicated 3 minutes until the mixture homogeneous that did not separate for at least two hours after sonicated. A typical preparation contained  $3.3 \times 10^{-3}$  M of phospholipid and  $3.3 \times 10^{-3}$  M of cholesterol in 1 litre of phosphate-buffered saline and 3 liters of solvent. PDG-SSL were sealed and sterilized. The size of the vesicles was determined by a dynamic light scattering technique. When PG/PC/CHOL were 1:4:5, diameter of liposomes was 20-50 nm (range). PDG-SSL was very stabilized in at least nine months.

So far, many articles reported drug-containing liposomes. However, liposomes are not stabilized enough. Therefore it is difficult to be used for pharmaceutical industry. In accordance with this invention, PDG-SSL is very stabilized in at least nine months. Therefore PDG-SSL can be used in industry. PDG-SSL can enhance targeting and improve pharmaceutical activity of PDG. In fact, PDG-SSL is safer than PDG. Also it is very important that all lipids are extracted from soybean. Therefore it is very safe for human being. In fact, many articles reported lipids, which used for drug-containing liposomes, are syntheses by organic chemistry. However synthetic lipids have some side effects, therefore, according to this invention, methods of preparing of PDG-SSL and phospholipids, which extracted from soybean, are very safe.

#### **Example 7 The effect of LX and PDG on control of oncogenes**

Human myeloblastic leukemic cell (ML-1) had been described previously (8). Cells were maintained in suspension culture in RPMI 1640 medium supplemented with 7.5% heat-inactivated FBS. Cells growth and viability were assayed by hemocytometer using trypan-blue dye exclusion. RNA was isolated by the CsCl gradient modification. RNA pellets were washed twice by reprecipitation in ethanol and quantitated by absorbency at 260nm. RNA analyzed by

electrophoresis of 15 µg of RNA through 1.2% agarose formaldehyde gels followed by northern blot transfer to nitrocellulose.

Single-standard uniformly labeled DNA probes were prepared. Probe of c-myc was a 1.7 Kb  $\lambda$ -Eco RI restriction fragment containing the 3'exon region of human c-myc and probe of c-myb was 1.0 Kb myb-specific Bam HI fragment. Probes for n-ras contained DNA fragments using a modification of the PCR technique. Probes for myb, myc and n-ras were isolated by electrolution. The isolated fragments were labeled to high specific activity with [ $\alpha^{32}$ P]-dCTP (3000 ci/mmol). Prehybridization of the filter was performed. The hybridization mixer contained 50,000 cpm of probe. The probes were hybridized at 58° C in 15 mM NaCl, 1.5 nM sodium citrate for 3 hours. After hybridization, they were exposed to XAR-5 film. Oncogene expression was quantitated by densitometer scanning of the autoradiography.

The effect of LX on oncogene was determined. The results are summarized in the tables as below.

Table 1. The effect of LX and PDG concentration on inhibition of oncogenes

LX concentration (ng/ml)	Inhibition (%)		
	c-myb RNA	c-myc RNA	RAS
Saline	0	0	0
LX (10)	25.0 $\pm$ 2.8*	24.2 $\pm$ 2.8*	15.2 $\pm$ 2.0*
LX (50)	68.5 $\pm$ 7.8**	78.0 $\pm$ 8.5**	28.2 $\pm$ 3.8**
LX (50) + PDG (50)	92.2 $\pm$ 12.5***	90.2 $\pm$ 10.5**	45.8 $\pm$ 6.2

nM = 10<sup>-9</sup> M; \*P<0.05, \*\*P<0.01, \*\*\*P<0.001 compared with LX

This study clearly indicated that LX and LX + PDG could significantly inhibit oncogenes of cancer cells. Cancers would be suitable targeted for gene-directed therapy and the present study has been directed toward the suppression of oncogene activity in cancers. Cellular oncogenes encode proteins have important function in differentiation of cancer cells. The principal functions of c-myc are the induction of proliferation and the inhibition of terminal differentiation in many cells. Over-expression of myc commonly occurs in a wide range of tumors.

#### **Example 8 The effect of LX and PDG on decreasing of tyrosine kinase**

In general, very low levels of TK are expressed in normal cells and high levels of TK are expressed in cancer cells. Many evidences have been accumulated that the dysfunction of cellular

oncogenes is a cause of human cancers. Therefore, a drug, which inhibits the activity of TK, can provide a new way to overcome cancer.

## Methods

**Cells.** L1210 and P388 cells were grown at 37°C on medium RPMI-1640 without antibiotics and supplemented with 10% horse serum. Cultures were diluted daily to  $1 \times 10^5$  cells/ml with fresh growth medium. From a culture initiated with cells from ascitic fluid obtained from a mouse 5 days after implantation with in vivo-passage leukemia, a stock of ampoule containing  $10^7$  cells/ml in growth medium plus 10% dimethyl sulfoxide was frozen and stored in liquid nitrogen.

Cultures were started from the frozen stock and were passage for no more than 1 month.

L1210 and P388 cells were grown at 37°C on medium RPMI-1640 supplemented with 10% calf serum, 10,000 unit/ml of Penicillin and 10,000 unit/ml of Streptomycin.  $1 \times 10^6$ /ml cells were placed in culture with different concentrations of LX. Then the cell suspension was incubated at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>-95% air for the indicated time. Reactions were terminated by addition of 3 ml of cold Earle's buffer. Cells were lysed, precipitated with 10% trichloroacetic acid (TCA) and filtered onto glass fiber filters. The filters were washed with phosphate-buffered saline and placed in scintillation vials, and radioactive emissions were counted.

H-60 leukemia cells were plated at a density of  $5 \times 10^5$  cells in 60-mm dish, and divided control and treatments groups for incubation 24 hours at 37°C with 5% CO<sub>2</sub>. The cells were collected and washed twice with phosphate-buffered saline, and resuspended at density of  $10^6$  cells/ml in 5mM HEPES buffer (pH 7.4). The cells were then resuspended in 1 ml of buffer containing 5mM HEPES (pH 7.6), 1mM MgCl<sub>2</sub> and 1mM EDTA, then placed on ice bath. The cell membrane was disrupted by ultra sound and centrifuged at 1000 x g for 10 minutes. The supernatant was ultra centrifuged at 30,000 X g for 30 minutes at 4°C. The pellet was resuspended in 0.3 ml of buffer containing 25mM HEOES, centrifuged at 12,000 x g for 5 minutes. The resulting supernatant was used for TK assay. Content of protein was determined. 10 µg of protein placed in 20mM HEPES (pH 7.6), 15mM MgCl<sub>2</sub>, 10mM ZnCl<sub>2</sub> and 5% (v/v) Nonidet P-40. After 5 minutes incubation at 25°C, the reaction was initiated by the addition of 25 µM [ $\gamma^{32}$ P] ATP (3 ci/mmol). After 10 minutes, the reaction was stopped by the addition of 20mM cold ATP. 50 µl of the mixtures were



spotted on glass microfiber filter discs and washed three times with cold trichloroacetic acid (TCA), contained 10mM sodium pyrophosphate. Air dried. Radioactivity was determined by liquid scintillation spectrometry. The net TK activity was determined after correcting for endogenous TK activity.

The present study clearly demonstrated that LX and LX + PDG significantly reduced in TK activity.

Table 2. Effect of LX and PDG on TK activity of HL-60 leukemia cells

Group	% of control activity
Control	100
LX (10ng/ml)	8.0
LX (50ng/ml)	32.5
LX (50ng/ml) + PDG (50ng/ml)	94.2

Value represents the mean of two experiments each done in duplicate; the range was less than 5% of the mean.

#### Example 9 LX and PDG controls differentiation of cancer cells

Human promyelocytic leukemia cells (HL-60) were grown in RPMI Medium 1640 supplemented with 10% (v/v) heat-inactivated FBS (56° C for 30 min) at 37° C in a humidified 95% air/5% CO<sub>2</sub> atmosphere. Cells were seeded at a level of  $2 \times 10^5$  cells/ml. Cells were allowed to attain a maximum density of  $1.2 \times 10^6$  cells/ml before being passed by dilution into fresh medium to a concentration of  $2 \times 10^5$  cells/ml.

Differentiation was induced in HL-60 cells by treatment with 0.1 µg/ml ( $10 \times 10^{-7}$  g/ml) of LX or PDG. Cell differentiation was measured by the ability of cells to reduce NBT.

Table 3. Induced differentiation of LX and PDG on cancer cells

Treatment	NBT%*
None	0
LX	$68.5 \pm 7.2$
LX + PDG	$72.8 \pm 8.5$

\*NBT represents the percentage of cells capable of reducing NBT.

The results of Table 4 indicate that LX and LX + PDG markedly induced differentiation of human cancer cells. Increasing NBT% means that cancer cells are differentiation.

### Example 10 Effects of LX and PDG on tumor cells proliferation

Human tumor cell lines: Hela leukemia HL-60, malignant melanocarcinoma B16, oral epidermoid carcinoma (KB), lung carcinoma (A549), breast carcinoma MCF-7, adenocarcinoma of stomach.

Animal tumor cell lines: Walker carcinoma, LLC-WRC-256, malignant melanoma (RMMI 1846), 3T3, and S-180 sarcoma (CCRF-180). All lines were routinely cultured in the RPMI1640 medium supplemented 20% fetal calf serum. The experiment was carried out in 96 microplate, each well had  $5 \times 10^5$  cells and given desired concentration of  $1 \mu\text{g/ml}$  ( $1 \times 10^{-6} \text{ g/ml}$ ) drug. Then the plate was incubated at  $37^\circ \text{C}$  in an atmosphere of humidified air enriched with 5 percent carbon dioxide for 72 hours. Concentration of LX or PDG is 50ng/ml.

Inhibition percent rate of tumor cell proliferation was obtained according to the bellow formula.

$$\text{Inhibition percent rate} = \frac{\text{Control} - \text{Test}}{\text{Control}} \times 100\%$$

LX inhibited tumor cells growth significantly. Percent rates of inhibition were all more than 70% in all cancer cells by LX.

Table 4. Effect of LX and PDG on inhibiting growth cancer cells

Cell line	Inhibition (%)	
	LX	LX + PDG
Control	---	---
Human cells		
HL-60	$78.5 \pm 8.2$	$82.8 \pm 9.0$
Hela	$70.7 \pm 8.0$	$84.0 \pm 9.8$
KB	$73.8 \pm 8.9$	$80.8 \pm 7.8$

### Example 11 Effect of LX and PDG on apoptosis of cancer cells

Human promyelocytic leukemia cells (HL-60) were grown in RPMI Medium 1640 supplemented with 10% (v/v) heat-inactivated FBS ( $56^\circ \text{C}$  for 30 min) at  $37^\circ \text{C}$  in a humidified 95% air/5%  $\text{CO}_2$  atmosphere. Cells were seeded at a level of  $2 \times 10^5$  cells/ml. Cells were allowed to attain

a maximum density of  $1.2 \times 10^6$  cells/ml before being passed by dilution into fresh medium to a concentration of  $2 \times 10^5$  cells/ml.

Apoptosis determined by the following process: Cell pellets containing  $5 \times 10^6$  cells were fixed with 2.5% glutaraldehyde in cacodylate buffer (pH 7.4), dehydrated through graded alcohol, and infiltrated with LX-112 epoxy resin. After overnight polymerization at  $60^\circ\text{C}$  1- $\mu\text{m}$  sections were cut with glass knives using a LKB Nova microtome. The sections were stained with 1% toluidine blue and coverslipped. In addition, experimental examples were stained with May-Grunwald-giemsa stain for the demonstration of apoptosis.

DNA electrophoresis: Untreated and treated HL-60 cells collected by centrifugation, washed in phosphate buffered saline and resuspended at a concentration of  $5 \times 10^6$  cells and 0.1% RNase A. The mixture was incubated at  $37^\circ\text{C}$  for 30 min and then incubated for an additional 30 min at  $37^\circ\text{C}$  with 1ml protease K. Buffer was added and 25  $\mu\text{l}$  of the tube content transferred to the Horizontal 1.5% agarose gel electrophoresis was performed at 2 V/cm. DNA in gels visualized under UV light after staining with ethidium Bromide (5 $\mu\text{g/ml}$ ).

DNA fragmentation assays: DNA cleavage was performed, quantitation of fractional solubilized DNA by diphenylamine assay and the percentage of cells harboring fragmented DNA determined by in labeling techniques. For the diphenylamine assay,  $5 \times 10^6$  cells were lysed in 0.5 mL lysis buffer (5 mmol/L Tris-HCl, 20 mmol/L DTA, and 0.5% Triton X-100, pH 8.0) at  $4^\circ\text{C}$ . Lysates were centrifuged (15,000g) for separation of high molecular weight DNA (pellet) and DNA cleavage products (supernatant). DNA was precipitated with 0.5N perchloric acid and quantitated using diphenylamine reagent.

The cell cycle distribution was determined 4 hours after addition of drug and represents mean  $\pm$  SD of 5 independent experiments.

Table 5 Effect of LX on apoptosis of cancer cells

Drug Concentration ( $\mu\text{g}$ )	Apoptosis (%)
Control	0
LX (50)	$70.8 \pm 8.2$
LX (50) + PDG (50)	$80.8 \pm 8.5$

Data of table 5 indicated that LX and LX + PDG could significantly induce apoptosis.

### Example 11 The effect of LX and PDG on the growth of animal transplanted tumor

Experimental procedure:

Male mice, weight 20 - 22g, were used in the experiment.  $1 \times 10^7$  tumor cells were injected to mouse and LX injected intraperitoneally began second day. All mice were sacrificed on the 12th days, isolated the tumor and weighed and calculated the inhibition rate of tumor weight.

Results:

The effect of LX on the growth of animal transplanted tumor as illustrated by the table 6. LX 20 mg/kg could inhibit the growth of S180, ECS, HCS, ARS, U-14 and L615 transplanted tumor.

Table 6. Inhibition rate (%) of transplanted tumor

Transplanted tumor	Inhibition (%)	
	LX (50ng/ml)	LX (50ng/ml) + PDG (50ng/ml)
Control	---	---
L1210	$74 \pm 9.0$	$87.0 \pm 9.8$
Lewis	$72 \pm 8.8$	$89.0 \pm 10.1$
S180	$70 \pm 9.0$	$83.0 \pm 9.8$
Walker 256	$75 \pm 8.0$	$92.5 \pm 10.8$

Data of table 6 indicated that LX and LX + PDG could significantly inhibit growth of tumor.

### Example 12 LX and PDG inhibited tumor incidence in vivo

The capacity of tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) to induce tumor incidence was recognized.

Methods:

Every group had 20 mice. For treatment group, each mouse was gave LX and PDG by injection at dose of 20mg/kg daily For control group, each mouse was gave same volume of physiological saline. Three days later, mice were gave 10  $\mu$ mol NNK (in 0.1ml saline) by i.p. injection. Sixteen weeks after these treatments the mice were killed and pulmonary adenomas were counted. The statistical significance of bioassay data was determined by student's test.

Table 7. Effects of LX and PDS on NNK-induced lung tumorigenesis

Group	Tumor incidence (%)	P
Control	100	---
LX	$28.2 \pm 5.0$	$<0.01$
LX + PDG	$20.0 \pm 3.0$	$<0.01$

Data of table 7 indicated that LX and LX + PDG have a significant tumor incidence.

### Example 13 The effect of LX and PDG on phagocytosis of peritoneal macrophage in mice

Added 0.02 ml of 5% washed chick red blood cell suspension to 0.5 ml of the peritoneal exudate, shaken gently to mix and incubated at 37°C for 5 minutes. Dipped two coverslips, closed to each other, in the above mixture and incubate for 30 minutes for the migration of the macrophages along the coverslips then fixed and stained with Sharma stain. Examined microscopically for: Phagocytic rate -- number of macrophages with phagocytized chick red blood cells per 100 macrophages counted.

Method of animal model was regular. The drugs were injected subcutaneously daily for 7 days.

Table 8. Effect of LX and PDG on immune function

Group	Number of samples	Phagocytic rate (x±SD, %)
Control	10	$34.10 \pm 1.45$
CY	10	$7.52 \pm 0.46$
CY + LX	10	$25.6 \pm 3.20^*$
CY + LX + PDG	10	$30.8 \pm 5.60^*$

\*P<0.01 compared with treatment group

The counting result showed that LX could increase the phagocytic rate. It suggested that LX and PDG could improve the nonspecial immune function in mice. And LX + PDG has more highly increasing immune function.

### Example 14 The effect of LX and PDG on the subpopulation of lymphocyte

As same as in Example 13, injected CY as the immune inhibitor for each mouse with one time daily and 7 times totally. The analysis of the subpopulation of T lymphocyte was carried out using fluorescein labeled monoclonal antibody of mouse CD<sub>8</sub> and CD<sub>4</sub> and determined cells by flow cytometer. The results are listed in Table 9.

Table 9. The effect of LX and PDG on the subpopulation of T lymphocyte

Group	Cases	CD <sub>8</sub>	CD <sub>4</sub>
Control	10	25.4	38.7
LX	10	19.0	20.2
LX + PA	10	33.0	36.8

Table 9 indicated that LX and LX + PDG could strongly increase immune function.

#### Example 15 Safety of composition

1. The acute LD<sub>50</sub> of LX was found to be 340mg/kg injection in abdominal cavity in mice. And LD<sub>50</sub> of LX-SSL was 721mg/kg. LD<sub>50</sub> of polysaccharide of Dang Gui for injection with abdominal in mice was higher than 1g/kg. They are very safe.
2. Each dose for an adult was 50-500 mg. Using 50 kg as the average weight of an adult the dosage of LX, LX-SSL, PDG and PDG-SSL were 1-10 mg/kg, and they are very safe.
3. As to subacute toxicity tests, a dosage corresponding to 50 times the clinical dose of LX was administered continually for two months, and no side effects had been observed. The electrocardiograms and functions of liver and the kidney had not been effect and no injuries whatever had been observed in the tissue slices of the heart, liver, spleen, lungs, kidneys and adrenal.

#### Example 16 Assay and identification of botanical drug

In the present study, we examined LX by chromatography with cambowax column. Temperature was 135°C. Methy salicylate was used as internal standard. The ration of peak arrears was linear function with concentration when between 0.5 to 1.5mg/ml. Data of LX are listed as below.

$$y = 1.32x - 0.03,$$

$$\gamma = 0.9999.$$

Molecular form: C<sub>15</sub>H<sub>24</sub>

Molecular weight: 204

Mp: 114~118°C

[α]<sup>16</sup>: -15°

IRv<sup>KBr</sup> cm<sup>-1</sup>: 3090, 2975, 2860, 1642, 1440, 1375, 1002, 910, 888 ( C=CH<sub>2</sub>);

PMR (CCl<sub>4</sub>)  $\delta$ : 0.97 (3H, s), 1.7 (6H, s), 4.4~5.6 (6H, m), 5.7 (1H, dd, = OH<sub>2</sub>);

MSm/e (%): 204 (M<sup>+</sup>), 147 (33), 121 (41), 107 (54), 93 (89), 81 (100), 79 (44), 68 (74), 67 (52), 55 (41), 53 (33), 41(52).

#### Example 17 Pharmacokinetics of LX

Distribution of LX (100mg/kg) after injection to rate was determined. The results are listed below table.

Table 10. Pharmacokinetics and disposition of LX

Organ	Concentration/ $\mu$ g/g		
	15 min	30 min	2 h
Heart	250 $\pm$ 32	15.8 $\pm$ 2.0	7.2 $\pm$ 1.0
Liver	125.5 $\pm$ 25	7.0 $\pm$ 0.8	0.7 $\pm$ 0.08
Spleen	150 $\pm$ 21	28 $\pm$ 3	2.0 $\pm$ 0.3
Lung	99 $\pm$ 10	10 $\pm$ 2.1	3.5 $\pm$ 0.4
Kidney	205 $\pm$ 31	50 $\pm$ 6.0	12 $\pm$ 2.0
Muscle	50 $\pm$ 6.1	9.0 $\pm$ 1.2	3.5 $\pm$ 0.4
Stomach	68 $\pm$ 7.0	8.9 $\pm$ 0.9	1.8 $\pm$ 0.2
Plasma	41 $\pm$ 5.0	6.5 $\pm$ 0.8	1.0 $\pm$ 0.2

The above data indicated that absorption and elimination of LX was rapid. LX, therefore, is a safe botanical drug.

The preparation of drugs, which can be accomplished by the extraction methods, set forth above or any conventional methods for extracting the active principles from the plants. The novelty of the present invention resides in the mixture of the active principles in the specified proportions to produce drugs, and in the preparation of dosage units in pharmaceutically acceptable dosage form. The term "pharmaceutically acceptable dosage form" as used hereinabove includes any suitable vehicle for the administration of medications known in the pharmaceutical art, including, by way of example, capsules, tablets, syrups, elixirs, and solutions for parenteral injection with specified ranges of drugs concentration.

In addition, the present invention provides novel methods for treating and preventing a variety of cancer conditions and control cancer cells with produced safe pharmaceutical agent.

It will thus be shown that there are provided compositions and methods which achieve the various objects of the invention and which are well adapted to meet the conditions of practical use.

As various possible embodiments might be made of the above invention, and as various changes might be made in the embodiments set forth above, it is to be understood that all matters herein described are to be interpreted as illustrative and not in a limiting sense.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000